

WHICH VACCINE DO I NEED?

Tetanus, Diphtheria, Pertussis (Tdap)

- You are entering the 6th grade
- You are between the ages of 10-12
- You are under the age of 64, never received Tdap before, and are due for your next Tetanus & Diphtheria (TD) booster (*needed roughly every 10 years*)
- You are under the age of 64 and are in close contact with infants under the age of 12 months (new mothers, fathers, daycare workers, grandparents, babysitters etc.)
- You got a severe cut or burn, protection after a wound

Meningococcal Vaccine

- You are entering the 6th grade
- You are between the ages of 11-12
- You are entering college for the first time and did not receive the vaccine earlier
- You are going back to college and will be living on campus and never got the vaccine previously
- You are traveling to a part of the world where the disease is common
- You were exposed to a Meningitis outbreak

Human Papillomavirus Vaccine (HPV)

- Males and Females between the ages of 9-26, regardless of whether you are sexually active or not, to prevent cervical cancer and other types of cancer as well as to prevent the most common sexually transmitted virus
- Recommended age for females 11 or 12 years of age
- Vaccine is a 3 dose series

Pneumococcal Vaccine

- You are above the age of 65
- Children under the age of 2, especially if they attend daycare
- You are between the ages of 2-64 and have one of the following:
 - a long-term health problem
 - a disease or condition that lowers your body's resistance to infection
 - you are taking a drug or treatment that lowers your body's resistance to infection
- You are between the ages of 19-64 and are a smoker or have asthma



